



#### **PHYSICAL PROPERTIES**

Appearance: Yellow Liquid

Specific Gravity: 1.0

**Odor:** Pleasant

Ph (Conc.): 6-7

Flash Point: >201°F (>93.9°C)

#### **PACKAGING**

1, 2.5 & 5 Gallon



**Q-64** 

# **DISINFECTANT CLEANER & DEODORIZER**

## **ABOUT THIS PRODUCT**

Q-64 Disinfectant Cleaner and Deodorizer is a phosphate-free, pH neutral formulation designed to provide effective cleaning, deodorizing, and disinfection for hospitals, nursing homes, schools, food processing plants, supermarkets, food service establishments, office buildings, manufacturing facilities, lodging establishments, retail business, athletic/recreation facilities and convention centers where housekeeping is of prime importance in controlling cross-contamination from treated surfaces. This product is a one step neutral disinfectant that is effective against a broad spectrum of bacteria, is virucidal\* (including HIV-1, HIV-2, HBV and HCV) and inhibits the growth of mold and mildew and their odors when used as directed. This product when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel, tables, chairs, desks, bed frames, walls, cabinets, doorknobs, garbage cans, glazed ceramic tile, plastic surfaces, play equipment, exercise equipment, exercise and wrestling mats, shower stalls, bathtubs, toilets and urinals.

#### **DIRECTIONS**

Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-11 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

One-step cleaning and disinfection of toilet bowls and urinals: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes) and flush.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a 3 minute contact time,

## **PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

# DANGER

KEEP OUT OF REACH OF CHILDREN.

CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. May be fatal if absorbed through skin. Harmful if swallowed. Wear goggles or face shield, rubber gloves, and protective clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

### STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

**PESTICIDE STORAGE**- Store in original container in a cool (temperature no lower than 50°F or higher than 120°F), dry and locked storage area inaccessible to children.

**PESTICIDE DISPOSAL-** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL-** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. After rinsing container, offer empty metal drums or plastic containers for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

RINSING INSTRUCTIONS- Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.