



Sanitizer L

LAUNDRY BACTERIOSTAT, SANITIZER, DISINFECTANT, *VIRUCIDE



PHYSICAL PROPERTIES

Appearance: Clear Liquid

Specific Gravity: 0.985

Odor: Characteristic

Ph (Conc.): 7-8

Flash Point: >201°F (93.9°C) estimated.

PACKAGING

4x1 gal

EPA REG. NO. 1839-155-74357

EPA EST. NO. 74357-MO-01

ABOUT THIS PRODUCT

Sanitizer L is designed specifically for hospitals, nursing homes, hotels, lodging establishments, schools, athletic/recreational facilities/tanning spas, food processing plants, dairies, food service establishments, restaurants, bars, transportation terminals, automotive garages, animal quarters, poultry and turkey farms, kennels and institutions where disinfection, sanitization, and deodorization is of prime importance. When used as directed, this product is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, sink tops, bathtubs, cabinets, tables, chairs, tanning beds and telephones.

Sanitizer L is a complete, chemically balanced disinfectant/sanitizer that provides clear use-solutions even when diluted in hard water.

DIRECTIONS

See the container label for complete directions for use.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical and semi-critical medical device surfaces.

DISINFECTION - To disinfect inanimate, hard, non-porous surfaces apply use-solution with mop, cloth, sponge, low pressure coarse sprayer or hand-pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

For low pressure coarse sprayer and hand pump trigger sprayer applications, spray 6-8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray.

Note: for spray applications, cover or remove all food products.

GENERAL DISINFECTION - Add 1.5 ounces of this product per 5 gallons of water for disinfection against *Staphylococcus aureus* (ATCC 6538), *Salmonella choleraesuis* (ATCC 10708), *Listeria monocytogenes* (ATCC 35152), and *Yersinia enterocolitica* (ATCC 23715).

HOSPITAL DISINFECTION - Add 1.75 ounces of this product per 5 gallons of water for disinfection against *Pseudomonas aeruginosa* (ATCC 15442) and Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), as well as the organisms listed under the GENERAL DISINFECTION and *VIRUCIDAL ACTIVITY sections.

KILLS HIV-1, HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

CUI Solutions, Inc.

1841 VERNON ST. | N. KANSAS CITY, MO 64116 | 816-471-3602

FAX: 816-474-3302 | WWW.CUISOLUTIONS.COM