



# Sani-T

## HARD SURFACE SANITIZER



### PHYSICAL PROPERTIES

- Appearance:** Colorless to red Liquid
- Specific Gravity:** 0.99
- Odor:** Characteristic
- Ph (Conc.):** 7-8
- Flash Point:** >201°F (93.9°C) estimated.

### PACKAGING

4X1 gal, 5 gal

**EPA REG. NO. 1839-86-74357**

**EPA EST. NO. 74357-MO-01**

### ABOUT THIS PRODUCT

Sani-T Hard Surface Sanitizer is designed for use in food processing plants, food service establishments, restaurants, bars, hospitals, nursing homes, hotels, schools, homes, dairies where, sanitization is of prime importance. Sani-T Hard Surface Sanitizer is for use as a sanitizer on dishes, glassware, utensils and food processing equipment at 200-400 ppm active quaternary without potable water rinse. Sani-T Hard Surface Sanitizer may be used on these washable hard, non-porous surfaces: food preparation and storage areas, dishes, glassware, silverware, cooking utensils, plastic and other non-porous cutting boards, coolers, ice chests, kitchen equipment, food processors, blenders, cutlery, utensils, countertops, stovetops, sinks, appliances, refrigerators, garbage cans, display equipment, high chairs, drinking fountains.

### DIRECTIONS

**Sanitization:** Recommended for use in restaurants, dairies, food processing plants, and bars. When used as directed this product is an effective sanitizer at an active quaternary concentration of 200-400 ppm when diluted in water up to 650 ppm hardness (CaCO<sub>3</sub>) in public eating establishments, dairies and food processing plants against Escherichia coli, Escherichia coli O157:H7, Staphylococcus aureus, Yersinia enterocolitica, Listeria monocytogenes, Salmonella typhi, Shigella sonnei, Vibrio cholera, Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), and Campylobacter jejuni.

Remove all gross food particles and soil from areas, which are to be sanitized, with a good detergent, pre-flush, pre-soak or pre-scrape treatment. Rinse with a potable water rinse.

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.